

## SUPPLEMENTARY INFORMATION

### Cell-Free miR-27a, a Potential Diagnostic and Prognostic Biomarker for Gastric Cancer

Jong-Lyul Park<sup>1</sup>, Mirang Kim<sup>1,2</sup>, Kyu-Sang Song<sup>3</sup>, Seon-Young Kim<sup>1,2</sup>,  
Yong Sung Kim<sup>1,2\*</sup>

<sup>1</sup>Epigenome Research Center, Genome Institute, KRIBB, Daejeon 34141, Korea,

<sup>2</sup>Department of Functional Genomics, Korea University of Science and Technology, Daejeon 34141, Korea,

<sup>3</sup>Department of Pathology, Chungnam National University College of Medicine, Daejeon 35015, Korea

**Supplementary Table 1.** Clinical characteristics of the plasma samples from 35 patients with gastric cancer and 35 healthy controls

Characteristics	Gastric cancer (n = 35)	Healthy control (n = 35)
Age (yr)	51.80 ± 11.88	48.94 ± 11.86
Sex		
Male	18 (51.4)	18 (51.4)
Female	17 (48.5)	17 (48.5)
Stage		-
I	16 (45.7)	-
II	8 (22.8)	-
III	9 (25.7)	-
IV	1 (2.8)	-
Not specified	1 (2.8)	-
Type		
Intestinal type	9 (25.7)	-
Diffused type	18 (51.4)	-
Mixed type	7 (20)	-
Not specified	1 (2.8)	-

Values are presented as mean ± standard deviation or number (%).